

Please replace the paragraph from lines 10-29 on page 40 with:

a2 Individual colonies are selected, cut with *AsuII* and *BamHI*, and the size of the *AsuII/BamHI* fragment determined. Colonies in which both the primer 1 and primer 2 sequences are correctly inserted are further amplified, an cut with *AsuII* and *BamHI* to produce the gene activation construct

FOET50-4964260  
cgaagcgaggcagccagcgagggagagagcgagcgggagcgagccgagcgaggaaATCGAAGGTTCGAATCCTTC  
CCCCACCACCATCACTTTCAAAAGTCCGAAAGAATCTGCTCCCTGCTTGTGTGTTGG  
AGGTCGCTGAGTAGTGCGCGAGTAAAATTTAAGCTACAACAAGGCAAGGCTTGACC  
GACAATTGCATGAAGAATCTGCTTAGGGTTAGGCGTTTTGCGCTGCTTCGCGATGTA  
CGGGCCAGATATACGCGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATT  
ACGGGGTCATTAGTTCATAGCCCATATATGGAGTTCGCGGTTACATAACTTACGGTA  
AATGGCCCGCCTGGCTGACCGCCCAACGACCCCCGCCCATTGACGTCAATAATGAC  
GTATGTTCCCATAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGACTA  
TTTACGGTAAACTGCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTACGCC  
CCCTATTGACGTCAATGACGGTAAATGGCCCGCCTGGCATTATGCCCAGTACATGAC  
CTTATGGGACTTTTCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATG  
GTGATGCGGTTTTTGGCAGTACATCAATGGGCGTGGATAGCGGTTTGA CTACGGGGA  
TTTCCAAGTCTCCACCCCATTTGACGTCAATGGGAGTTTGT TTTGGCACCAAAATCAA  
CGGGACTTTCCAAAATGTCGTAACAACTCCGCCCCATTGACGCAAATGGGCGGTAG  
GCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCTCTGGCTAACTAGAGAACCCA  
CTGCTTACTGGCTTATCGAAATTAATACGACTCACTATAGGGAGACCCAAGCTTGGT  
ACCGAGCTCGGATCgatctgggaaagcgcaagagagagcgcacacgcacacacccgcccgcgcactcgg (SEQ ID  
No. 25)

Please replace the first paragraph on page 53 with:

In an illustrative embodiment, the *ptc* therapeutic can be an antisense construct for inhibiting the expression of *patched*, e.g., to mimic the inhibition of *patched* by *hedgehog*.

a3 Exemplary antisense constructs include:

5'-GTCCTGGCGCCGCGCCGCGCCGTCGCC (SEQ ID No. 26)

5'-TTCCGATGACCGGCCTTTCGCGGTGA (SEQ ID No. 27)

5'-GTGCACGGAAAGGTGCAGGCCACACT (SEQ ID No. 28)

*The replacement paragraphs presented above incorporate changes as indicated by the marked-up versions below.*

In an exemplary embodiment, portions of the 5' flanking region of the human Shh gene are amplified using primers which add restriction sites, to generate the following fragments

5'-gcgcgcttcgaaGCGAGGCAGCCAGCGAGGGAGAGAGCGAGCGGGCGAGCCGGAGC-GAGGAAatcgatgcgcgc (primer 1) (SEQ ID No. 23)

5'-gcgcgcagatctGGGAAAGCGCAAGAGAGAGCGCACACGCACACACCCGCCGCGCG-CACTCGggatccgcgcgc (primer 2) (SEQ ID No. 24)

Individual colonies are selected, cut with AsuII and BamHI, and the size of the AsuII/BamHI fragment determined. Colonies in which both the primer 1 and primer 2 sequences are correctly inserted are further amplified, an cut with AsuII and BamHI to produce the gene activation construct

cgaagcgaggcagccagcgagggagagagcgagcgggagcgagcggagcgaggaaATCGAAGGTTCGAATCCTTC  
CCCCACCACCATCACTTTCAAAGTCCGAAAGAATCTGCTCCCTGCTTGTGTGTTGG  
AGGTCGCTGAGTAGTGCGCGAGTAAATTTAAGCTACAACAAGGCAAGGCTTGACC  
GACAATTGCATGAAGAATCTGCTTAGGGTTAGGCGTTTTGCGCTGCTTCGCGATGTA  
CGGGCCAGATATACGCGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATT  
ACGGGGTCATTAGTTCATAGCCCATATATGGAGTTCGCGTTACATAACTTACGGTA  
AATGGCCCGCCTGGCTGACCGCCCAACGACCCCGCCATTGACGTCAATAATGAC  
GTATGTTCCCATAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGACTA  
TTTACGGTAAACTGCCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTACGCC  
CCCTATTGACGTCAATGACGGTAAATGGCCCGCCTGGCATTATGCCCAGTACATGAC  
CTTATGGGACTTTTCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATG  
GTGATGCGGTTTTTGGCAGTACATCAATGGGCGTGGATAGCGGTTTGA CTACGGGGA  
TTTCCAAGTCTCCACCCCATGACGTCAATGGGAGTTTGT TTTGGCACCAAAATCAA  
CGGGACTTTCCAAAATGTCGTAACA ACTCCGCCCCATTGACGCAAATGGGCGGTAG  
GCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCTCTGGCTAACTAGAGAACCCA  
CTGCTTACTGGCTTATCGAAATTAATACGACTCACTATAGGGAGACCCAAGCTTGGT  
ACCGAGCTCGGATCgatctgggaaagcgcaagagagagcgcacacgcacacccgcgcgcactcgg (SEQ ID  
No. 25)